

No. of Printed Pages : 11

## **MPH-002/MPH-003**

**M. SC. (PHYSICS) (MSCPH)**

**Term-End Examination**

**June, 2025**

**MPH-002 : CLASSICAL MECHANICS-I**

**&**

**MPH-003 : ELECTROMAGNETIC THEORY**

*Time : 3 Hours*

*Maximum Marks : 50*

***Instructions :***

- 1. Students registered for both MPH-002 and MPH-003 courses should answer both the question papers in two separate answer books entering their enrolment number, course code and course title clearly on both the answer books.*
- 2. Students who have registered for any of the MPH-002 or MPH-003 should answer the relevant question paper after entering their enrolment number, course code and course title on the answer book.*

**Part-A****MPH-002 : CLASSICAL MECHANICS—I***Time :  $1\frac{1}{2}$  Hours**Maximum Marks 25*

---

**Note :** *All questions are compulsory. However, internal choices are given. Marks for each question are indicated against it. You may use a calculator. Symbols have their usual meanings.*

---

---

1. Attempt any *one* part : 5×1=5

- (a) A particle of mass  $m$  moving with velocity  $\vec{v}$ , undergoes an elastic collision with a particle of mass  $3m$  which is initially at rest. The kinetic energy of the particle ( $m$ ) after collision is one-fourth of its kinetic energy before

the collision. Calculate the angles of the direction of motion of the two particles after the collision using conservation of linear momentum.

- (b) Write down the equation of constraint for two particles connected by a massless rod of length  $L$ . The system is moving in  $X$ - $Y$ -plane such that the velocity of the centre of the rod is along its length.

2. Attempt any *one* part :  $5 \times 1 = 5$

- (a) A cylinder of radius  $R$  and mass  $m$  rolls without slipping down an inclined plane of length,  $s$  making an angle  $\theta$  with the

horizontal. Obtain the Lagrangian and determine the equation of motion if the moment of inertia of the cylinder is

$$\frac{mR^2}{2}.$$

- (b) A spherical pendulum of mass  $m$  and length  $A$ , has the Lagrangian :

$$L = \frac{1}{2}mA^2(\dot{\theta}^2 + \dot{\phi}^2 \sin^2 \theta) + mgA \cos \theta$$

Determine the generalized momenta and energy function.

3. Attempt any *one* part : 5×1=5

- (a) Derive the Euler-Lagrange equation of motion for a simple pendulum with a bob of mass  $2m$  which is tied to a string of length  $L_0$ , where  $L_0$  is a constant.

- (b) Obtain the effective potential for a particle of mass  $m$  with angular momentum  $l$ , undergoing motion along a spiral path of the form  $r = Ae^{b\theta}$  when the total energy is zero.  $A$  and  $b$  are positive constants.

4. Attempt any *one* part : 10×1=10

- (a) Consider two coupled pendulums each of mass  $m$  and length  $l$ . Write down the  $V$  and  $T$  matrices for the system. It is given that  $k$  is the spring constant of the spring connecting the bobs. 10
- (b) A particle of mass  $m$  moves in a central force field described by the potential

$$V(r) = \frac{-k}{r} \exp\left(\frac{-r}{a}\right), \text{ where } k \text{ and } a \text{ are}$$

positive constants.

(i) Determine the force. 2

(ii) If the motion of the particle is restricted to a plane  $(r, \dot{\theta})$ , write the Lagrangian and obtain the equation of motion for  $r$ . 7

(iii) Obtain the effective potential. 1

## Part—B

## MPH-003 : ELECTROMAGNETIC THEORY

*Time :  $1\frac{1}{2}$  Hours*

*Maximum Marks : 25*

---

**Note :** *All questions are compulsory. Marks for each question are indicated against it. Symbols have their usual meanings. You may use a calculator.*

---

1. Answer any *three* parts : 3×5=15
- (a) A coaxial cable consists of a thin inner solid copper wire and an outer sheath of braided copper wire. The linear charge density of the inner wire is  $-\lambda$  and that of the outer wire is  $\lambda$ . Using Gauss'

law, determine the electric fields at a point : 3+2

(i) in the region inside the inner wire,  
and

(ii) in the region between the wires.

(b) Write Poisson's equation for the charge

density of a wire given by  $\rho(z) = \frac{2\rho_0 x}{L}$ ,

where  $L$  is the length of the wire.

Determine the electric potential and electric field due to the wire for the following boundary conditions : 5

$$V(x) = 0 \text{ at } x = 0$$

and  $\frac{\partial V(x)}{\partial x} = 0 \text{ at } x = L.$



- (c) Assuming that the electrons in the atoms of dielectric materials are bound to their respective nuclei by harmonic forces, show that the polarisation is given by :

$$\vec{P} = \frac{Nq^2}{m\omega_0^2} \vec{E}$$

where  $\omega_0$  is the natural frequency of oscillations of electron about the nucleus.

5

- (d) Calculate the magnetic field  $\vec{B}$  and the magnetising field  $\vec{H}$  at (i) a point 90 mm from a long straight wire carrying a current of 10 A and (ii) the centre of a 1500 turn solenoid which is

0.20 m long and carries a current 1.5 A  
flowing in it. (3+2)

[Take  $\mu_0 = 4\pi \times 10^{-7} \text{ Hm}^{-1}$ ]

- (e) Derive the wave equation for magnetic field  $\vec{B}$  from Maxwell's equations for vacuum. 5

2. Answer any *one* part : 1×10=10

- (a) (i) Show that for a linear and isotropic dielectric kept in an electric field, the surface bound charge density is given by : 5

$$\sigma_b = \vec{P} \cdot \hat{n}$$

- (ii) Show that for non-uniform polarisation of a dielectric placed in an electric field, the bound charge density is given by : 5

$$\rho_b = -\nabla \cdot \vec{P}$$

- (b) Explain the concept of magnetic vector potential. Show that the magnetic vector potential due to an infinite straight wire carrying current  $I$ , at a distance  $r$  from the wire is given by :

$$\vec{A}(r) = -\frac{\mu_0 I}{2\pi} \ln(r) \hat{z}$$

where  $\hat{z}$  denotes the direction of current. 2+8

× × × × ×